



PEST OFF!

Deadly effective

RODENT BLOCKS

NET WEIGHT 10kg



HARMFUL SUBSTANCE - ECOTOXIC

KEEP OUT OF REACH OF CHILDREN - READ ATTACHED LABEL CAREFULLY BEFORE USE

For use in bait stations for controlling rodents. Contains 0.02g/kg of brodifacoum in the form of a cereal composite block bait. Store in original container, tightly closed and away from feed or foodstuffs.

Keep out of reach of children and domestic animals.

Registered pursuant to the
ACVM Act 1997, No. V005099

Registered to & manufactured by:
Animal Control Products Ltd
408 Heads Road,
Whanganui 4501, NZ
www.pestoff.co.nz

Distributed by:

Bell Booth
Bell-Booth Ltd
15 Tiki Place,
Palmerston North, NZ
FREEPHONE 0800 80 90 91
www.bell-booth.co.nz

HSNO CLASSES 6.9B, 9.1D
HSNO Approval Code: HRC000004
For safety data sheet visit
www.pestoff.co.nz

Approved as a Maintenance
Compound Type C and
MPI recognised for use
in dairy premises



DIRECTIONS FOR USE

Mice: Place 50-100 pellets or a single block into a Pestoff Mouse Bait Station (or similar bait station) and place bait stations in areas used by mice but out of reach of non-target animals, children and wildlife. Mice are attracted to sheltered, warm areas near or under refrigerators, stoves, hot water cylinders and cupboards. Check bait stations and replenish product daily until baits cease to be eaten. Repeat periodically to check any re-infestation.

Rats: Place 1-2 cups of pellets or 4 to 8 blocks into a Pestoff Dead Rat Café (or similar bait station) and place bait stations in areas used by rats but out of reach of non-target animals, children and wildlife. Rats have poor eyesight and use fixed objects, such as runways along walls and banks for navigation. Place bait stations on these runways or near burrows. Check bait stations and replenish pellets or blocks daily until baits cease to be eaten. Repeat periodically to check any re-infestation.

Residue Warning: Brodifacoum, the active ingredient, is a potent second generation anticoagulant poison which can accumulate particularly in the liver, kidneys and fat of poisoned animals. Do not feed rodents killed with brodifacoum baits to dogs or cats.

Legal obligations: These products must only be used in bait stations. If product is applied in areas to which the public has right of access, it is a legal requirement that signs must be posted to notify the public that this product has been applied in the area. Signs must remain for a period of 12 months after product has been removed or destroyed. This product must only be used as specified in the label.

Shelf life: The shelf life of this product may vary according to the suitability of storage conditions. Any product held after the expiry date shown on container should be disposed of according to label directions.

Livestock: It is important to prevent access to blocks and baits by domestic livestock and pets. Stock must be kept off the treatment area until the product has been removed or destroyed. Dogs and cats are at risk from scavenging poisoned animal carcasses and pet owners in the immediate vicinity should be notified of this risk. Collect poisoned animal carcasses where practicable for burning or burying at least 600mm below ground, otherwise limit access to the treatment area until poisoned animal carcasses are unlikely to be eaten or to contain residues.

Maintenance compound status: Pestoff Rodent Blocks are approved as a maintenance compound Type C. They are also MPI recognised for use in dairy processing under the conditions that they:

1. must not be used in areas accessible to food producing animals,
2. must be used in a suitable bait station and
3. may only be used in manufacturing environments in accordance with a documented pest programme.

Conditions of sale: As no control can be exercised over the methods or conditions under which this product is used, no responsibility or claim, other than those required by statute, will be accepted for any damage or injury whatsoever arising from the storage, handling, application, use or disposal of this product.



Distributed by Bell-Booth Limited
15 Tiki Place, Palmerston North, New Zealand
FREEPHONE 0800 80 90 91

For safety data sheet go to:
<http://www.pestoff.co.nz>
Registered pursuant to the ACVM Act 1997
No. V005099 (Pestoff Rodent Blocks)
No. V005137 (Pestoff Rodent Pellets)
See: <http://www.foodsafety.govt.nz>
for registration conditions.

Registered to and manufactured by
Animal Control Products Ltd
408 Heads Road, Whanganui, New Zealand

PESTOFF® RODENT PELLETS & PESTOFF® RODENT BLOCKS
Harmful. Keep out of reach of children. Ecotoxic
HSNO CLASSES 6.9B, 9.1D
Baits in pellet or block form for use in bait stations for controlling rodents.
Active ingredient: 0.02g/kg brodifacoum

PRECAUTIONS

Handling: Repeated oral exposure may cause toxin to accumulate in internal organs and may affect the clotting ability of the blood. Wear impervious gloves when handling the product or open containers. Do not eat, drink or smoke when using the product or handling open containers. Remove protective clothing and wash hands and exposed skin thoroughly before meals and after any contact. Do not breathe the dust from these products.

Toxic to terrestrial vertebrates: Take measures to prevent domestic animals and pets being exposed to the toxin either through eating the product or through eating the carcasses of poisoned animals. Apply these products in bait stations only.

Storage: Do not store in direct sunlight. Store in original container, tightly closed and away from feed or foodstuffs. Keep out of reach of children and domestic animals.

EMERGENCY MANAGEMENT

Symptoms of Poisoning: No symptoms may be apparent for several days if poisoning has occurred. Symptoms may include pale gums, passing of blood in urine or faeces, and the appearance of bruising.

First Aid: If poisoning is suspected DO NOT induce vomiting. Call a doctor or emergency physician at your nearest hospital. For further advice contact National Poisons Centre 0800 POISONS (Phone 0800 764 766).

Treatment of domestic animals accidentally poisoned: Vitamin K1 is an effective antidote against accidental poisoning of domestic animals by brodifacoum (the active ingredient). Veterinarians are familiar with the information on Vitamin K1 therapy and should be consulted.

Spillage: In the event of a spill, isolate the spill area and take all practicable steps to manage any harmful effects of a spillage including preventing product from entering streams or waterways. Scoop spilled product into secure containers. Recover any undamaged product for later use by placing in appropriately labelled containers and dispose of spoiled product as directed below. Use a broom to collect fine material and wash down the spill area with copious water only after all spilled product has been removed.

Disposal: Product which is surplus or spoiled should be disposed of by burying with other organic material on the active tip face of an appropriately managed landfill or buried within the biologically active layer of soil elsewhere within a secure area. Ensure that a good covering of earth is applied immediately to prevent access by scavenging birds. Alternatively, burn unwanted product in a suitably constructed and appropriately located incinerator (if permitted) and bury any residues as above. Treating the product through a sewage oxidation facility or other chemical treatment facility is also an acceptable means of disposing of unwanted bait material. Dispose of empty container at a landfill. Do not use the empty container for any other purpose.

WJS:ACP JULY 2015

PLEASE READ BEFORE USE

PEST OFF!
Deadly effective

Open along here

IMPORTANT
PRODUCT INFORMATION

GLUE HERE

GLUE HERE



Red - Pantone 200 C



Green - Pantone 7724 C

Size 65mm high x 305mm wide